



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ture which it would have were it immediately raised from water under the pressure which it has acquired by compression or expansion. The only cause of the restoration of vapour to the liquid form is the abstraction of heat from it; and this cause will be equally operative, whatever may be the state of the vapour with respect to density: but compression alone, without such abstraction of redundant heat, can never convert any portion of vapour into a liquid. In accordance with these views, the author regards the permanent gases as vapours, containing a large quantity of redundant heat.

A paper was also read, entitled, "On the Secretion and Uses of the Bile." By B. Phillips, Esq. Communicated by W. G. Maton, M.D. V.P.R.S.

The object of the author of this paper is to establish the three following propositions; viz.

1°. That the principles of the bile pre-exist in the blood, and that the function of the liver is to separate from the blood a certain proportion of this material.

2°. That bile may be secreted as well from arterial as from venous blood.

3°. That chyle may be formed in the absence of bile.

In support of the first proposition the author adduces the analogy of other secretions, the suppression of which is followed by the appearance in the blood of the peculiar animal product which characterizes that secretion, as has been proved in the case of the urine by Prévost and Dumas, and afterwards by Vauquelin, Serullas, and Magendie. The author has confirmed the conclusion to which these physiologists have arrived, by some experiments of his own, in which, instead of extirpating the kidneys, he contented himself with tying the venal vessels. He relates two cases in which the vena portæ and hepatic artery were tied, and bile was found both in the urine and the blood.

The author, after quoting several authorities in support of his second proposition, that the liver can secrete bile, although the vena portæ be obstructed, relates two experiments which he made on dogs, by tying the vena portæ at the part before it arrives at the transverse fissure of the liver: in both cases that organ continued to secrete bile, though the quantity was small. In another dog, he tied the hepatic artery, with the effect of producing fatal peritonitis, but without any apparent change in the biliary secretion.

The arguments adduced by the author in favour of the opinion that chyle may be formed when no bile is present in the intestine, are derived from the accounts given by various authors of cases in which the ductus communis had been rendered impervious by the pressure of neighbouring tumours. In confirmation of this result, he made experiments on four dogs, and found chyle in the thoracic duct after he had tied the ductus communis close to the duodenum.

The author concludes, from these and other facts, that the secretion of bile is intended to serve some other purpose than that of contributing to the formation of chyle.